

Product number **T009**  
Revision number **RN3.0**

---

<b>Product Name</b>	Human keratinocyte transglutaminase (TG1, recombinantly produced in <i>E. coli</i> )
<b>Synonym</b>	TG1, TGase 1, keratinocyte protein-glutamine- $\gamma$ -glutamyltransferase
<b>Source</b>	Recombinant, produced in <i>E. coli</i>
<b>Quantity</b>	150 $\mu$ g
<b>Description</b>	<p>His<sub>6</sub>-rhTG1 is based on the TGM1-allele from I.M.A.G.E.-clone IRAKp961M1628 isolated from human skin squamous cell carcinoma. It is N-terminally fused to a hexahistidine-tag resulting in the encoded N-terminal amino acid sequence MHHHHHHMDGPR.</p> <p>His<sub>6</sub>-rhTG1 is purified by IMAC to more than 90 % purity.</p>
<b>Molecular Weight</b>	90 kDa
<b>Activity</b>	> 2,500 U/mg [Activity is determined by measuring the rate of fluorescence enhancement after His <sub>6</sub> -rhTG1-catalyzed monodansylcadaverine-incorporation into N,N-dimethylated casein according to Lorand et al., Anal. Biochem. 44 (221-231) see fig.2]. 1 U is defined as the increase in fluorescence intensity of 1 a.u./min (measured on a Cary eclipse fluorescence spectrophotometer, Varian; $\lambda_{ex}$ = 332 nm, $\lambda_{em}$ = 500 nm; band filter = 5 nm; detector strength = 600 V; temperature = 37 °C, assay volume = 1 ml)].
<b>Activation</b>	Add 10 mM Ca <sup>2+</sup> to activate His <sub>6</sub> -rhTG1.
<b>Appearance</b>	White lyophilized solid.
<b>Application</b>	His <sub>6</sub> -rhTG1 catalyzes acyl transfer reactions from glutamin residues in proteins or peptides to primary amines, e. g. the formation of $\epsilon$ -( $\gamma$ -glutamyl) lysine bonds between proteins by transferring the acyl group of a peptide-bound glutamine residue to the primary amino group of a peptide-bound lysine residue. His <sub>6</sub> -rhTG1 may also be used for immunoprecipitation.
<b>Reagents</b>	The Transglutaminase is lyophilized from 50 mM Tris-HCl pH 8.0, 10 mM Glutathion.
<b>Reconstitution</b>	Add the volume of water specified in the certificate of analysis under aliquotation to the vial of lyophilized powder. Rotate vial gently until solid dissolves. After reconstitution the solution should be cooled on ice for short term storage.
<b>Storage</b>	Store at $\leq$ - 20°C. Store working aliquots at $\leq$ - 20°C. Avoid repeated freezing and thawing. <b><i>Delivery at ambient temperature is possible</i></b>
<b>Reference(s)</b>	Chrobok et al., PLoS One. 2018, 13:e0196433 Plank et al., J Invest Dermatol. 2018, pii: S0022-202X(18)32817-3 Fischer et al., J. Invest. Dermatol. 2013, 133:1170-7 Fukui et al., FEBS J. 2013, 280:1420-9 Schaertl et al., J. Biomol. Screen. 2010, 15:478-87 Yamane et al., FEBS J. 2010, 277:3564-74
<b>Related products</b>	T035 Human keratinocyte transglutaminase (TG1, recombinantly produced in insect cells) A018 Polyclonal antibody to human keratinocyte transglutaminase A029 FITC-labelled polyclonal antibody to human TG1 T101 1,3,4,5-Tetramethyl-2[(2-oxo-propyl)thio] imidazolium chloride T036 Transglutaminase Assay Kit, fluorescent, Casein, Dansylcadaverine
<b>Release date</b>	23 December 2021

# Product Data Sheet



Product number **T009**  
Revision number RN3.0

---

**NOTE** INTENDED FOR RESEARCH USE ONLY, NOT FOR USE IN HUMAN, THERAPEUTIC OR DIAGNOSTIC APPLICATIONS.